arXiv:1401.7083v2 [math.GR] 8 Mar 2015

Extensions of locally compact abelian, torsion-free groups by compact torsion abelian groups

Hossein SAHLEH and Ali Akbar ALIJANI

Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Guilan, Rasht, Iran
sahlehg@gmail.com

Abstract. Let X be a compact torsion abelian group. In this paper, we show that an extension of F_p by X splits where F_p is the p-adic number group and p a prime number. Also, we show that an extension of a torsion-free, non-divisible LCA group by X is not split.

Introduction

Throughout, all groups are Hausdorff abelian topological groups and will be written additively. Let $\pounds$ denote the category of locally compact abelian (LCA) groups with continuous homomorphisms as morphisms. The Pontrjagin dual of a group G is denoted by $\hat{G}$. A morphism is called proper if it is open onto its image and a short exact sequence $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ in $\pounds$ is said to be proper exact if ϕ and ψ are proper morphisms. In this case the sequence is called an extension of A by C (in $\pounds$). Following [5], we let $Ext(C, A)$ denote the (discrete) group of extensions of A by C. The splitting problem in LCA groups is finding conditions on A and C under which $Ext(C, A) = 0$. In [3, 4, 6] the splitting problem is studied. We have studied the splitting problem in the category of divisible, LCA groups [9]. By using the splitting problem, we determined the LCA groups G such that the maximal torsion subgroup of G is closed [10]. Let X be a compact torsion group. In [4, Theorem 1], it is proved that if G is a divisible LCA group, then $Ext(X, G) = 0$. However, the suggested proof in [4] appears to be incomplete as it uses the incorrect Proposition 8 of [3]. In [9], we proved that if G is a divisible, $\sigma-$compact group, then $Ext(X, G) = 0$. Let P be the set of all prime numbers, J_p, the p-adic integer group and F_p, the p-adic number group which is the minimal divisible extension of J_p for every $p \in P$ [7]. By [7, 25.33], a divisible, torsion-free LCA group G has the form $G \cong \mathbb{R}^n \bigoplus A \bigoplus M \bigoplus E$, where A is a discrete, torsion-free, divisible group, M a compact connected torsion-free group and E, the minimal divisible extension of $\prod_{p\in P} J_p^{n_p}$ where n_p is a cardinal number for every $p \in P$. In general, $E \neq \prod_{p\in P} F_p^{n_p}$. Let I be a finite subset of P such that

$$n_p = \begin{cases} 0 & \text{if } p \notin I \\ 1 & \text{if } p \in I \end{cases}$$

2010 *Mathematics Subject Classification.* 20K35, 22B05.
Key words and phrases. locally compact abelian group, extension, divisible group, torsion-free group.

Then the minimal divisible extension of $\prod_{p\in I} J_p^{n_p}$ is $\prod_{p\in I} F_p^{n_p}$. In this paper, we show that if $G \cong \mathbb{R}^n \bigoplus A \bigoplus M \bigoplus \prod_{p\in I} F_p^{n_p}$, then $Ext(X, G) = 0$ (see Theorem 1.6).

The additive topological group of real numbers is denoted by $\mathbb{R}$, $\mathbb{Q}$ is the group of rationales with discrete topology and $\mathbb{Z}$ is the group of integers. The topological isomorphism will be denote by " $\cong$ ". For more on locally compact abelian groups see [7].

1. Extensions of a torsion-free LCA group by a compact torsion abelian group

Lemma 1.1. *Let $X \in \pounds$ and p a prime number. Then $nExt(X, F_p) = Ext(X, F_p)$ for every positive integer n.*

Proof. Let n be a positive integer and $f : F_p \to F_p$, $f(x) = nx$ for all $x \in F_p$. By [1, Lemma 2], f is open. So f is a proper morphism. Consider the exact sequence $0 \to Kerf \to F_p \xrightarrow{f} F_p \to 0$. By Corollary 2.10 of [5], we have the exact sequence

$$\to Ext(X, Kerf) \to Ext(X, F_p) \xrightarrow{f_*} Ext(X, F_p) \to 0 \tag{1.1}$$

Since $f_*(Ext(X, F_p)) = nExt(X, F_p)$, it follows from sequence (1.1) that $nExt(X, F_p) = Ext(X, F_p)$. □

Lemma 1.2. *Let X be a compact torsion group. Then $Ext(X, F_p) = 0$.*

Proof. F_p is a totally disconnected group. So, by Theorem 24.30 of [7], F_p contains a compact open subgroup K. Now we have the following exact sequence

$$... \to Ext(X, K) \to Ext(X, F_p) \to Ext(X, F_p/K) \to 0 \tag{1.2}$$

Since F_p is divisible, so $Ext(X, F_p/K) = 0$ (see [5, Theorem 3.4]). Since X is compact and torsion, so by [7, Theorem 25.9], $nX = 0$ for some positive integer n. Hence, $nExt(X, K) = 0$ (see [8, Lemma 2.5]). Since (1.2) is exact, so $nExt(X, F_p) = 0$. Hence by Lemma 1.1, $Ext(X, F_p) = 0$. □

Remark 1.3. Let X be a group and $f : X \to X$,$f(x) = nx$ for all $x \in X$. If f is a topological isomorphism for every positive integer n, then X is a divisible, torsion-free group.

Theorem 1.4. *Let X be a compact group and p a prime number. Then $Ext(X, F_p)$ is a divisible, torsion-free group.*

Proof. Let n be a positive integer. Then the exact sequence $0 \to X \xrightarrow{\times n} X \to X/nX \to 0$ induces the following exact sequence

$$Ext(X/nX, F_p) \to Ext(X, F_p) \xrightarrow{\times n} Ext(X, F_p) \to 0$$

By Lemma 1.2, $Ext(X/nX, F_p) = 0$. So $Ext(X, F_p) \xrightarrow{\times n} Ext(X, F_p)$ is a topological isomorphism. Hence by Remark 1.3, $Ext(X, F_p)$ is a divisible, torsion-free group. □

Corollary 1.5. *Let $X \in \pounds$. Then $Ext(X, F_p)$ is a divisible, torsion-free group.*

Proof. Let $X \in \mathcal{L}$. By [7, Theorem 24.30], $X = \mathbb{R}^n \bigoplus H$ where H contains a compact open subgroup K. Consider the exact sequence

$$Ext(H/K, F_p) \to Ext(H, F_p) \to Ext(K, F_p) \to 0$$

Since H/K is a discrete group and F_p a divisible group, so $Ext(H/K, F_p) = 0$. Hence $Ext(H, F_p) \cong Ext(K, F_p)$. By Theorem 1.4, $Ext(K, F_p)$ is a divisible, torsion-free group. So $Ext(X, F_p)$ is a divisible, torsion-free group. □

Theorem 1.6. *Let X be a compact torsion group and $G \cong \mathbb{R}^n \bigoplus A \bigoplus M \bigoplus \prod_{p\in I} F_p^{n_p}$ where I is a finite subset of P defined as follows:*

$$n_p = \begin{cases} 0 & \text{if } p \notin I \\ 1 & \text{if } p \in I \end{cases}$$

Then $Ext(X, G) = 0$.

Proof. First recall that by [5, Theorem 2.13],

$$Ext(X, G) \cong Ext(X, A) \bigoplus Ext(X, M) \bigoplus \prod_{p\in I} Ext(X, F_p)$$

Since X is a totally disconnected group, so by [5, Theorem 3.4], $Ext(X, A) = 0$. Also $Ext(X, M) \cong Ext(\hat{M}, \hat{X})$. Since $\hat{X}$ is a discrete bounded group and $\hat{M}$ a discrete torsion-free group, so by [2, Theorem 27.5],$Ext(\hat{M}, \hat{X}) = 0$. By Lemma 1.2, $Ext(X, F_p) = 0$. Hence $Ext(X, G) = 0$. □

Lemma 1.7. *Let X be a compact torsion group. Then $Hom(X, \mathbb{Q}/\mathbb{Z}) \cong \hat{X}$.*

Proof. The exact sequence $0 \to \mathbb{Z} \to \mathbb{Q} \to \mathbb{Q}/\mathbb{Z}$ induces the following exact sequence

$$Hom(X, \mathbb{Q}) \to Hom(X, \mathbb{Q}/\mathbb{Z}) \to Ext(X, \mathbb{Z}) \to Ext(X, \mathbb{Q})$$

Since X is torsion and $\mathbb{Q}$ is torsion-free, so $Hom(X, \mathbb{Q}) = 0$. Also by [5, Theorem 3.4], $Ext(X, \mathbb{Q}) = 0$. Hence $Hom(X, \mathbb{Q}/\mathbb{Z}) \cong Ext(X, \mathbb{Z})$. By [5, Theorem 2.12 and Proposition 2.17], $Ext(X, \mathbb{Z}) \cong Ext(\hat{\mathbb{Z}}, \hat{X}) \cong \hat{X}$. So $Hom(X, \mathbb{Q}/\mathbb{Z}) \cong \hat{X}$. □

Theorem 1.8. *Let X be a compact torsion group and G a torsion-free, non-divisible group. Then $Ext(X, G) \neq 0$.*

Proof. Let G^* be the minimal divisible extension of G. By [7, A.13], G^* is a divisible, torsion-free group. Since X is torsion and G^* torsion-free, so $Hom(X, G^*) = 0$. By [5, Corollary 2.10], we have the following exact sequence

$$0 = Hom(X, G^*) \to Hom(X, G^*/G) \to Ext(X, G)$$

Since G^*/G is a discrete, torsion divisible group, so $Hom(X, G^*/G)$ containing a copy of $Hom(X, \mathbb{Q}/\mathbb{Z})$. Hence by Lemma 1.7, $Ext(X, G) \neq 0$. □

References

1. Armacost D. L. On pure subgroups of LCA groups. Trans. Amer. Math. Soc 1974;45: 414-418.
2. Fuchs L. Infinite Abelian Groups. New York: Academic Press, 1970.
3. Fulp R. O. Homological study of purity in locally compact groups. Proc. London Math. Soc 1970; 21: 501-512.
4. Fulp R. O. Splitting locally compact abelian groups. Michigan Math. J 1972;19: 47-55.
5. Fulp R. O, Griffith P. Extensions of locally compact abelian groups I. Trans. Amer. Math. Soc 1971;154: 341-356.

6. Fulp R. O, Griffith P. Extensions of locally compact abelian groups II. Trans. Amer. Math. Soc 1971;154: 357-363.
7. Hewitt E, Ross K. Abstract Harmonic Analysis. Berlin: Springer-Verlag, 1979.
8. Loth P. On t-pure and almost pure exact sequences of LCA groups. J. Group Theory 2006;9: 799-808.
9. Sahleh H, Alijani A. A. Splitting of extensions in the category of locally compact abelian groups. Int. J. Group Theory 2014;3:3: 39-45.
10. Sahleh H, Alijani A. A. The closed maximal torsion subgroup of a locally compact abelian group. JARPM 2014;6:3: 5-11.